

The SPICT™ is a guide to identifying people at risk of deteriorating health and dying. Assess these people for unmet supportive and palliative care needs.

Look for two or more general indicators of deteriorating health.

- Performance status is poor or deteriorating (the person is in bed or a chair for 50% or more of the day); reversibility is limited.
- Dependent on others for most care needs due to physical and/or mental health problems.
- Two or more unplanned hospital admissions in the past 6 months.
- Significant weight loss (5-10%) over the past 3-6 months, and/ or a low body mass index.
- Persistent, troublesome symptoms despite optimal treatment of underlying condition(s).
- Patient asks for supportive and palliative care, or treatment withdrawal.

Look for any clinical indicators of one or more advanced conditions

Cancer

Functional ability deteriorating due to progressive metastatic cancer.

Too frail for oncology treatment or treatment is for symptom control.

Dementia/ frailty

Unable to dress, walk or eat without help.

Eating and drinking less; swallowing difficulties.

Urinary and faecal incontinence.

No longer able to communicate using verbal language; little social interaction.

Fractured femur; multiple falls.

Recurrent febrile episodes or infections; aspiration pneumonia.

Neurological disease

Progressive deterioration in physical and/or cognitive function despite optimal therapy.

Speech problems with increasing difficulty communicating and/ or progressive swallowing difficulties.

Recurrent aspiration pneumonia; breathless or respiratory failure.

Heart/ vascular disease

NYHA Class III/IV heart failure, or extensive, untreatable coronary artery disease with:

- breathlessness or chest pain at rest or on minimal exertion.

Severe, inoperable peripheral vascular disease.

Respiratory disease

Severe chronic lung disease with:

- breathlessness at rest or on minimal exertion between exacerbations.

Needs long term oxygen therapy.

Has needed ventilation for respiratory failure or ventilation is contraindicated.

Kidney disease

Stage 4 or 5 chronic kidney disease (eGFR < 30ml/min) with deteriorating health.

Kidney failure complicating other life limiting conditions or treatments.

Stopping dialysis.

Liver disease

Advanced cirrhosis with one or more complications in past year:

- diuretic resistant ascites
- hepatic encephalopathy
- hepatorenal syndrome
- bacterial peritonitis
- recurrent variceal bleeds

Liver transplant is contraindicated.

Review supportive and palliative care and care planning

- Review current treatment and medication so the patient receives optimal care.
- Consider referral for specialist assessment if symptoms or needs are complex and difficult to manage.
- Agree current and future care goals, and a care plan with the patient and family.
- Plan ahead if the patient is at risk of loss of capacity.
- Record, communicate and coordinate the care plan.